



**Iron Infusion Request Form**

**SEND REFERRAL**  
Email: referrals@topdoc.au  
Fax: (02) 4096 1371

**MORE INFORMATION**  
1300 070 731

- Biologic Infusions
- Venesection
- Bisphosphonates
- + More Services

SCAN QR FOR ALL SERVICES & REFERRAL RESOURCES

**PATIENT DETAILS** Please fill in if no patient label available

SURNAME: .....

GIVEN NAMES: .....

DOB: ..... SEX:  M  F  Other

\*\*\*\*\* AFFIX PATIENT LABEL HERE \*\*\*\*\*

**TopDoc Clinic**  
Location:

PLEASE VISIT OUR WEBSITE FOR  
FULL LIST OF LOCATIONS

**Patient Clinical Details**

**Patient Support**

**CLINICAL INDICATION / DIAGNOSIS**

- Iron Deficiency
- Iron Deficiency Anaemia
- Other: .....

**MEDICAL HISTORY:**

- Nil Known
- Heart Failure
- Renal Failure
- Liver Disease
- Fluid Restriction
- Severe Asthma
- Pregnant → Gestation (weeks) .....
- Other: .....

**ALLERGIES / ADVERSE DRUG REACTIONS**

- Nil Known
  - Previous Infusion Reaction
  - Medicine / Other
- DESCRIBE REACTION

**CLINICAL PARAMETERS:** CAN LEAVE BLANK IF BLOODS ATTACHED

Height ..... Weight .....

CREAT ..... eGFR .....

Corr Ca ..... Ferritin .....

Hb ..... Hct .....

\*\*\*\*\* PLEASE ATTACH LATEST BLOOD RESULTS \*\*\*\*\*

- Booking Assistance Required:**  
Our nursing team will contact this patient to assist them in booking their appointment
- Phone no: .....
- Mobile Infusion Service Required:**  
Our outreach nursing team will provide the infusion at the patients home or residential aged care facility

- Interpreter Required:**  
LANGUAGE
- Aboriginal / Torres Strait Islander**

**Medical Procedure**

**Iron Infusion**

**MEDICATION / DOSE**

- Ferinject® 500mg (single dose)
- Ferinject® 1000mg (single dose)
- Ferinject® 1500mg (divided dose)
- Ferinject® 2000mg (divided dose)
- Monofer® 500mg (single dose)
- Monofer® 1000mg (single dose)
- Monofer® 1500mg (single dose)
- Monofer® 2000mg (divided dose)

**SIMPLIFIED DOSING TABLE**

Hb (g/L)	Weight 35-70 kg	Weight ≥70 kg
≥140	500 mg	500 mg
100-139	1,000 mg	1,500 mg
<100	1,500 mg	2,000 mg

Source: National Blood Authority, 2016; eMIMS Elite, 2024

**PROTOCOL**

- Standard Protocol:** Intravenous (Ferinject/Monofer) in 100 mL 0.9% NaCl, infused over 30 minutes.
- Special Protocol:** DILUENT/ DURATION/TREATMENT SCHEDULE

**Pre-medication (optional)**

**MEDICATION / DOSE**

- Loratadine 10mg PO
- Hydrocortisone 50mg / 100mg IV (circle one)
- Other: .....

**PLEASE NOTE**

Pre-medication is not routinely required but may be considered in patients with multiple drug allergies, asthma, or prior severe infusion reactions.

**Referrer Authorisation**

**DOCTOR / NP NAME:** .....

**PROVIDER NO:** .....

**PRACTICE NAME:** .....

**BY SIGNING, I CONFIRM THE FOLLOWING:**

I have explained the purpose, risks, and side effects of this treatment. The patient understands they may withdraw consent at any time.

The patient has no known contraindications and I have provided a **valid prescription** (eScripts preferred), along with **relevant blood results** (<4 weeks old) and supporting clinical documentation.

**DOCTOR / NP SIGNATURE:**

**DATE:**

.....

.....

DO NOT WRITE IN THIS BINDING MARGIN



1

**Book Your Appointment & Upload Your Referral**

5 min

2

**Complete Patient Registration Form**

3 min

3

**Attend Your Scheduled Appointment**


30 - 90 min

## Understanding Your IV Procedure

You have been prescribed an intravenous (IV) procedure. This involves placing a small, soft plastic tube called a cannula into a vein, usually in your arm or hand. The cannula allows fluids or medications to be delivered, or blood to be collected, in a safe and comfortable manner by an accredited nurse.

### What to Expect

- ▶ A tourniquet is applied to help locate a suitable vein.
- ▶ A needle is used briefly to insert the cannula. Once in place, the needle is retracted and only the soft plastic tube remains in the vein.
- ▶ A dressing is applied to secure the cannula in place.
- ▶ Your treatment will then begin. This may involve fluids or medication being slowly infused, or a prescribed amount of blood being collected, depending on the procedure prescribed.
- ▶ Once the procedure is complete, the cannula is removed, pressure is applied to the site to stop any bleeding, and a small bandage is placed over the area.



Example of IV Cannulation

## Medication Collection

If your procedure requires medication, you may either collect it from your local pharmacy or opt in to our **Medication Collection Service** through the **Patient Registration Form**. If you choose this option, we will arrange collection on your behalf and ensure your medication is ready for your appointment. Please contact us if you require any assistance.

## Getting Ready for Your Appointment

### Before You Arrive:

- ✔ Eat a light meal to avoid nausea or dizziness.
- ✔ Stay well hydrated unless advised otherwise
- ✔ Wear loose clothing with sleeves that roll up easily.
- ✔ Bring a jacket in case you feel cold.
- ✔ Please bring your Medicare card, any applicable concession cards, and a valid form of photo ID

### When You Arrive:

- ✔ Arrive 10 minutes early for check-in and preparation.
- ✔ Let staff know if you're unwell or if your health has changed.
- ✔ You will be asked to complete a pre-procedure assessment and consent form.
- ✔ A nurse will explain the procedure, ensure your comfort, insert the cannula, and monitor you throughout your procedure.

### After Your Procedure:

- ✔ Your IV procedure may take 30–90 minutes depending on your treatment.
- ✔ You will be asked to remain for up to 30 minutes after your procedure so we can ensure you are feeling well before you leave.
- ✔ Mild bruising or tenderness at the cannulation site is common and usually settles within a few days.
- ✔ Check the area daily for a week. If you notice discolouration, pain, swelling or discharge, contact us or your doctor.

### Health & Safety Notice

If you have a heart or kidney condition, please check with your doctor regarding fluid restrictions prior to your procedure. Please do not attend the clinic if you have cold or flu symptoms. Contact us to reschedule your appointment if you are feeling unwell.

Your doctor has recommended TopDoc Infusion Clinic for your IV procedure. While you are free to choose any licensed medical infusion provider, please note that not all clinics may offer the specific medications or services required for your care. We recommend discussing your options with your treating practitioner to ensure continuity, safety, and the best outcome for your treatment.

DO NOT WRITE IN THIS BINDING MARGIN

PATIENT INFORMATION FOR INTRAVENOUS (IV) PROCEDURE